

INDEX TO VOLUME 61

AUTHORS

- Allen, Larry G.: *see* Crane, Allen and Eisemann, 68-94
- Anderson, M. Eric and Gregor M. Cailliet: Occurrence of the rare North Pacific frostfish, *Benthodesmus elongatus pacificus*, Parin and Becker, 1970, in Monterey Bay, California, 149-152; *see* Cailliet and Anderson, 60-62
- Andreasen, James K.: Occurrence of the fathead minnow, *Pimephales promelas*, in Oregon, 155-156
- Bertram, Ronald C.: *see* Rempel and Bertram, 237-239
- Burton, Timothy S.: *see* Piekielek and Burton, 4-25
- Cailliet, Gregor M. and M. Eric Anderson: Occurrence of the prowfish, *Zaprora silenus* Jordan, 1896 in Monterey Bay, California, 60-62; *see* Anderson and Cailliet, 149-152
- Campbell, Gail and Robson A. Collins: The age and growth of the Pacific bonito, *Sarda chiliensis*, in the eastern North Pacific, 181-200
- Clark, Patrick, James Nybakken and Lawrence Laurent: Aspects of the life history of *Tresus nuttallii* in Elkhorn Slough, 215-227
- Collins, Robson A.: *see* Campbell and Collins, 181-200
- Courtois, Louis A.: Blood and serum analysis of adult striped bass captured in the Sacramento River, 245-246
- Crane, Jules M., Larry G. Allen and Connie Eisemann: Growth rate, distribution, and population density of the northern quahog *Mercenaria mercenaria* in Long Beach, California, 68-94
- Crase, Frederick T.: *see* DeHaven, Crase and Woronecki, 166-180
- DeHaven, Richard W., Frederick T. Crase and Paul P. Woronecki: Breeding status of the tricolored blackbird, 1969-1972, 166-180
- deWit, Leray A.: Changes in the species composition of sharks in south San Francisco Bay, 106-111
- Dinnel, Paul A.: *see* Quirolo and Dinnel, 156-157
- Duffy, John M.: A range extension and two new California size records for mollusks, 152-155
- Eisemann, Connie: *see* Crane, Allen and Eisemann, 68-94
- Espinosa, F. A. Jr.: *see* Paulson and Espinosa, 200-214
- Eldridge, Maxwell B.: Early larvae of the diamond turbot, *Hyposopsetta guttulata*, 26-34
- Fry, Donald H.: Measuring salmon, an old and unfamiliar method, 247
- Gall, G. A. E.: *see* Gold and Gall, 248-250
- Gold, J. R. and G. A. E. Gall: Further record of the Little Kern golden trout, *Salmo aguabonita whitei* in the Little Kern River Basin, California, 248-250
- Green, Roger E.: A preliminary list of fishes collected from Richardson Bay, California 1972-1973, 104-106
- Hauser, William J.: An unusually fast growth rate for *Tilapia zillii*, 54-56
- Hensley, Gary H. and F. M. Nahhas: Parasites of fishes from the Sacramento-San Joaquin Delta, California, 201-208
- Hobson, Edmund S.: First California record of the serranid fish *Anthias gordensis* Wade, 111-112
- Knaggs, Eric H., John S. Sunada and Robert N. Lea: Notes on some fishes collected off the outer coast of Baja California, 56-59
- Knight, Allen W.: *see* Kost and Knight, 35-46
- Kost, Angela L. Blado and Allen W. Knight: The food of *Ncomysis mercedis* Holmes in the Sacramento-San Joaquin Estuary, 35-46
- Laurent, Lawrence: *see* Clark, Nybakken and Laurent, 215-227
- Lea, Robert N.: *see* Knaggs, Sunada and Lea, 56-59
- Mathews, C. P.: Note on the ecology of the ratfish, *Hydrolagus collei*, in the Gulf of California, 47-53
- McCullough, Dale R.: Modification of the Clover deer trap, 242-244
- Mearns, Alan J.: *Pociliopsis gracilis* (Heckel), newly introduced pociliid fish in California, 251-253
- Odenweller, Dan B.: An unusual aggregation of bat rays, *Myliobatis californica* Gill, 159
- Nahhas, F. M.: *see* Hensley and Nahhas, 201-208
- Nybakken, James: *see* Clark, Nybakken and Laurent, 215-227

- Paulson, Larry J. and F. A. Espinosa, Jr.: Fish trapping: A new method of evaluating fish species composition in limnetic areas of reservoirs, 209-214
- Piekielek, William and Timothy S. Burton: A black bear population study in northern California, 4-25
- Quirolo, Lawrence F. and Paul A. Dinnel: Latitudinal range extensions for yellow and spotted snake eels (genus *Ophichthus*), 156-157
- Rawstron, Robert R.: Mortality and growth rates, cost, and relative contribution of two different sizes of silver salmon stocked in Lake Berryessa, California, in 1972, 127-132
- Rempel, Ronald D. and Ronald C. Bertram: The Stewart modified corral trap, 237-239
- Russo, Ronald A.: Notes on the external parasites of California inshore sharks, 228-232; Observations on the food habits of leopard sharks (*Triakis semifasciata*) and brown smoothhounds (*Mustelus henlei*), 95-103
- Schott, Jack W.: Otter trawl cod-end escapement experiments for California halibut, 82-103
- Spratt, Jerome D.: Anomalous otoliths from the northern anchovy, *Engraulis mordax*, 235-236; Growth rate of the northern anchovy, *Engraulis mordax*, in southern California waters, calculated from otoliths, 116-126
- Sunada, John S.: Age and length composition of northern anchovies, *Engraulis mordax*, in the 1972-73 season, California anchovy reduction fishery, 133-143; see Knaggs, Sunada and Lea, 56-59
- Talent, Carline L.: see Talent and Talent, 233-234
- Talent, Larry G.: A parakeet auklet, *Cyclorhynchus psittacula*, from Monterey Bay, California, 158; Pugheadedness in the longspine combfish, *Zaniolepis latipinnis*, from Monterey Bay, California, 160-162
- Talent, Larry G. and Carline L. Talent: An extrauterine fetus in the Steller sea lion, *Eumetopias jubata*, 233-234
- Thomas, Ronald D.: The status of Rocky Mountain elk in Kern County, 1974, 239-241
- White, Robert L.: Polymorphism in populations of *Sceloporus occidentalis* in Santa Barbara County, California, 253-254
- Wilbur, Sanford R.: California condor plumage and molt as field study aids, 144-148
- Wintersteen, John: Occurrence and depth range extension of the yellow snake eel, *Ophichthus zophochir*, off southern California, 157-158
- Woronecki, Paul P.: see DeHaven, Crase and Woronecki, 166-180

SCIENTIFIC NAMES

- | | |
|--|---|
| <i>Acanthomysis sculpta</i> : 45 | <i>Benthodesmus simonyi</i> : 149 |
| <i>Accipiter gentilis</i> : 145 | <i>Benthodesmus</i> sp.: 149 |
| <i>Acer macrophyllum</i> : 6 | <i>Brassica campestris</i> : 172 |
| <i>Acipenser medirostris</i> : 206 | <i>Bothriocephalus claviceps</i> : 201; 205 |
| <i>Acipenseridae</i> : 206 | <i>Branchellion lobata</i> : 230-231 |
| <i>Aegypius monachus</i> : 146 | <i>Branchellion</i> sp.: 230 |
| <i>Agonopsis sterletus</i> : 58 | <i>Callianasa californiensis</i> : 99 |
| <i>Alloglossidium corti</i> : 201; 204 | <i>Callianasa</i> sp.: 99, 102 |
| <i>Alosa sapidissima</i> : 201; 205-206 | <i>Cancer productus</i> : 99 |
| <i>Anchoa exiguus</i> : 57 | <i>Cancer</i> sp.: 99 |
| <i>Anoplopoma fimbria</i> : 58 | <i>Capillaria catenata</i> : 201; 205 |
| <i>Anthias gordenis</i> : 111 | <i>Carassius auratus</i> : 201, 205 |
| <i>Anthias sechurae</i> : 111 | <i>Carthamus tinctorius</i> : 172 |
| <i>Apolymetis biangulata</i> : 72 | <i>Catostomus rimitulus</i> : 155 |
| <i>Agelaius tricolor</i> : 166 | <i>Centaurea</i> sp.: 172 |
| <i>Aquila chrysaetos</i> : 146 | <i>Ceratium</i> sp.: 44 |
| <i>Arbutus menziesii</i> : 6 | <i>Cervus canadensis canadensis</i> : 239 |
| <i>Arctostaphylos patula</i> : 6 | <i>Cestoda</i> : 201, 204 |
| <i>Arctostaphylos viscida</i> : 6 | <i>Chacnobryttus gulosus</i> : 201; 204 |
| <i>Arculatala demissa</i> : 72 | <i>Chione fluctifraga</i> : 72-73 |
| <i>Arundo donax</i> : 172 | <i>Chione undatella</i> : 72 |
| <i>Atherinopsis californiensis</i> : 110 | <i>Chlorophyta</i> : 44 |
| <i>Atherinopsis</i> sp.: 101 | <i>Chromis punctipinnis</i> : 111 |
| <i>Atractolytceustus huronensis</i> : 201, 204-205 | <i>Ciona intestinalis</i> : 72 |
| <i>Benthodesmus elongatus pacificus</i> : 149-150 | <i>Citharichthys</i> sp.: 100 |
| | <i>Cleidodiscus pricei</i> : 201; 204 |
| | <i>Clevelandia ios</i> : 100 |

- Clinocardium nattalli*: 215
Clinostomum marginatum: 201, 204
Clupea harengus pallasi: 101
Clupeidae: 206
Contracacum brachyurum: 201, 206
Contracacum spiculigerum: 201, 206
Corallobothrium fimbriatum: 201, 205
Corallobothrium giganteum: 201, 205
Cornus nuttallii: 6
Coscinodiscus sp.: 40-42
Cottidae: 206
Cottus gulosus: 206
Crago franciscorum: 99
Crustacea: 201, 206
Cryptophyta: 44
Cyanophyta: 44
Cyclorhynchus psittacula: 158
Cyclotella sp.: 41
Cymatogaster aggregata: 100
Cynoscion nobilis: 106
Cyprinidae: 207
Cyprinodon macularius: 252
Cyprinus carpio: 201, 203-205, 252
Dactylogyrus extensus: 201, 204
Digena: 204
Dorosoma cepedianum: 209
Dorosoma petenense: 206, 209
Echthrogaleus coleoptratus: 229
Embiotocidae: 207
Engraulis mordax: 100, 116, 133, 182, 235
Enteromorpha intestinalis: 72
Enteromorpha sp.: 72, 79
Entosphenus tridentatus: 155
Epinephelus nireatus: 58
Eptatretus stoutii: 56
Eumetopias jubata: 233
Florimetis obesa: 72
Galeorhinus zyopterus: 110, 228
Gambusia affinis: 251
Gari californica: 153-154
Gymnogyps californianus: 144
Haliacetus leucocephalus: 146
Hemigrapsus nudus: 99
Hemigrapsus oregonensis: 99, 110
Hemigrapsus sp.: 99
Heterandria pleurospilus: 251
Hexanchus griseus: 57
Hirudinea: 201, 206
Hydrolagus collei: 47, 57
Hydrolagus sp.: 57
Hypsopsetta guttulata: 26, 33
Hypsopsetta sp.: 26, 33
Hysteroecarpus traskii: 207
Ictalurus catus: 201, 204-206
Ictalurus melas: 201, 205
Ictalurus nebulosus: 201, 204, 206
Ictalurus punctatus: 201, 204-205
Illinobdella alba: 206
Illinobdella elongata: 206
Illinobdella moorei: 201, 206
Illinobdella richardsoni: 206
Illinobdella sp.: 206
Keratella sp.: 44
Khavia iowensis: 201, 205
Lacistorhynchus sp.: 201, 205
Laevicardium elatum: 72
Lavinia exilicauda exilicauda: 207
Lepomis cyanellus: 205
Lepomis macrochirus: 201, 204-205
Lerhaea cyprinacea: 201, 206
Lerneopidae: 229
Lerneopoda scylica: 229
Libocedrus decurrens: 6
Lycodopsis pacifica: 158
Melosira sp.: 40-42
Mercenaria mercenaria: 68, 72-74, 78-80
Mercenaria sp.: 72-74, 78, 80
Metabronema salvelini: 201, 206
Microstomus pacificus: 59, 158
Molgula manhattenensis: 102
Monogenea: 204
Morone saxatilis: 35, 110, 201, 204-206, 245
Mustelus henlei: 95, 228
Mya arenaria: 217
Myliobatis californica: 100, 159
Mylopharodon conocephalus: 201, 206
Mysis relicta: 45
Mytilus edulis: 72
Myxobolus cyprinicola: 204
Myxobolus ellipsoides: 204
Myxobolus koi: 201, 204
Myxobolus sp.: 203-204
Myxobolus toyamai: 204
Nematoda: 201, 205
Neomysis integer: 45
Neomysis mercedis: 35, 40, 44
Neomysis sp.: 45
Notemigonus crysoleucas: 207
Notorhynchus maculatus: 100, 110, 228
Notropis lutrensis: 251
Octopus sp.: 102
Odocoileus virginianus: 242
Oncorhynchus kisutch: 127
Oncorhynchus tshawytscha: 6, 18, 207
Ophichthus sp.: 156
Ophichthus triserialis: 157
Ophichthus zophochir: 156-157
Orthodon microlepidotus: 201, 206
Pachygrapsus crassipes: 99
Pandaris bicolor: 228-229
Paralichthys californicus: 82
Paramysis arenosa: 45
Parophrys retulus: 58, 85
Pediastrum sp.: 44
Pelichnibothrium speciosum: 201, 205
Perissopus oblongatus: 228-229
Philometra carassii: 201, 205
Philometra sanguinea: 205
Philometra sp.: 205
Philometra triliabiata: 205
Physiculus rastrelliger: 57
Pinephales promelas: 155-156
Pinnipedia: 233
Pinus lambertiana: 6
Pinus ponderosa: 6
Pinus sabiniana: 6
Piscicolidae: 230
Platichthys stellatus: 207
Pleuronectidae: 207

- Pleuronichthys coenosus*: 33
Pleuronichthys decurrens: 26, 33
Pleuronichthys sp.: 26, 33
Pleuronichthys verticalis: 26, 33, 83
Pocilia latipinna: 252
Pociliopsis gracilis: 251-253
Pociliopsis mexicana: 251
Pociliopsis occidentalis: 251
Poecilistes pleurospilus: 251
Pogonichthys macrolepidotus: 201
Pogonichthys macrolepidotus: 206
Pomoxis nigromaculatus: 201, 206, 211
Porichthys notatus: 96, 100, 158
Posthodiplostomum minimum centrarchi:
 201, 204
Prionace glauca: 229
Protothaca staminea: 72-73, 153, 215
 Protozoa: 201
Psettichthys melanostictus: 85
Pseudotsuga menziesii: 6
Ptychocheilus grandis: 201, 206
Quercus garryana: 6
Quercus kelloggii: 6
Quercus wislizenii: 6
Raja binoculata: 229
Raja trachura: 106
Raphidascaris sp.: 201, 206
Rhinichthys osculus: 155
Rubus sp.: 172
Salix sp.: 171
Salmo aguabonita: 248
Salmo aguabonita aguabonita: 248, 250
Salmo aguabonita whitei: 248
Salmo gairdneri: 127, 155, 207, 248,
 250
 Salmonidae: 207
Sarda chilensis: 181
Sardinops sagax caeruleus: 116
Saridomus nuttalli (= *S. nuttallii*):
 153
Saridomus nuttallii (= *S. nuttalli*):
 215
Sceloporus occidentalis: 253-254
Sceloporus occidentalis biserialatus: 253
Sceloporus occidentalis bocourti:
 253-254
Sceloporus occidentalis longipes:
 253-254
Sceloporus occidentalis occidentalis: 253
Scenedesmus sp.: 44
Schistomysis spiralis: 45
Schizothaerus nuttalli: 101
Schizothaerus sp.: 101, 215
Scirpus sp.: 171
Scomber japonicus: 125, 182, 188
Scyllium sp.: 229
Semele decisa: 153-154
Spiroxya sp.: 201, 206
Squalus acanthias: 100, 228
Squatina californica: 230
Stereolepis gigas: 57
Styella plicata: 72
Sula bassana: 145
Symphurus atricauda: 158
Tabellaria sp.: 43
Tagelus californicus: 72
Terebra pedroana: 152, 154
Tilapia mossambica: 252
Tilapia sp.: 54, 252
Tilapia zillii: 54-55
Trachurus symmetricus: 182
Tresus capax: 215, 226
Tresus nuttalli (= *T. nuttallii*): 153
Tresus nuttallii (= *T. nuttalli*): 215,
 221, 226
Tresus sp.: 215, 219, 226
 Trematoda: 201
Triakis semifasciata: 95, 228
Triceratium sp.: 43
Typha sp.: 171
Ulva lobata: 72
Upogebia pugetensis: 99
Upogebia sp.: 99, 102
Urechis caupo: 101
Urechis sp.: 102
Ursus americanus: 4
Urtica sp.: 172
Vultur gryphus: 146
Xiphophorus helleri: 252
Xiphophorus variatus: 252
Xystreus liolepis: 83
Zaniolepis frenata: 58
Zaniolepis latipinnis: 58, 160
Zaprora silenus: 60
Zaprora sp.: 61
Zostera marina: 101

SUBJECT

- Age: of Pacific bonito in eastern North Pacific, 181-200
 Age composition: of northern anchovies in the 1972-73 season California anchovy reduction fishery, 133-143
 Analysis: of blood and serum of adult striped bass captured in Sacramento River, 245-246
 Anchovy, northern: age and length composition in the 1972-73 season California anchovy reduction fishery, 133-143; anomalous otoliths in, 235-236; growth rate in southern California waters, 116-126
 Anchovy reduction fishery: age and length composition of northern anchovies in, 1972-73, 133-143
 Areas, limnetic: evaluating fish species composition in reservoirs by fish trapping, 209-214
 Auklet, parakeet: found near Monterey Bay, 158
 Bass, striped: analysis of blood and serum of adults captured in Sacramento River, 245-246

- Bear, black: a population study in northern California, 4-25
- Blackbird, tricolored: breeding status, 1969-1972, 166-180
- Bonito, Pacific: age and growth in eastern North Pacific, 181-200
- Combfish, longspine: pugheadedness in, 160-162
- Components: of blood and serum from adult striped bass captured in Sacramento River, 245-246
- Condor, California: plumage and molt as field study aids, 144-148
- Contribution: of two different sizes of silver salmon stocked in Lake Berryessa, 127-132
- Cost: of two different sizes of silver salmon stock in Lake Berryessa, 127-132
- Deer: Trapping with modified Clover deer trap, 242-244; trapping with Stewart modified corral trap, 237-239
- Distribution: of northern quahog in Long Beach, California, 68-81
- Ecology: of ratfish in Gulf of California: 47-53
- Elk, Rocky Mountain: status in Kern County, 1974, 239-241
- Elkhorn Slough: site of study on life history of *Tresus nuttallii*, 215-227
- Evaluation: of fish species composition in limnetic areas of reservoirs with fish trapping, 209-214
- Fetus, extrauterine: in Steller sea lion, 233-234
- Fish, poeciliid: newly introduced in California, 251-253
- Fishes: collected from Richardson Bay, 1972-73, 104-106; collected off the outer coast of Baja California, 56-59
- Food: of *Neomysis mercedis* in the Sacramento-San Joaquin Estuary, 35-46
- Food habits: of leopard sharks, 95-103; of brown smoothhounds, 95-103
- Frostfish, North Pacific: occurrence in Monterey Bay, 149-152
- Growth: of Pacific bonito in eastern North Pacific, 181-200
- Growth rate: of northern anchovy in southern California waters, 116-126; of northern quahog in Long Beach, California, 68-81; of *Tilapia zillii*, 54-56
- Growth rates: of two different sizes of silver salmon stocked in Lake Berryessa, 127-132
- Gulf of California: site of study on the ecology of the ratfish, 47-53
- Halibut, California: otter trawl cod-end escapement experiments for, 82-94
- Kern County: status of Rocky Mountain elk, 1974, 239-241
- Lake Berryessa: site of mortality and growth rate study of two different sizes of stocked silver salmon, 127-132
- Larvae: of diamond turbot, 26-34
- Length composition: of northern anchovies in 1972-73 season California anchovy reduction fishery, 133-143
- Life history: of *Tresus nuttallii* in Elkhorn Slough, 215-227
- Little Kern River Basin: further record of Little Kern golden trout, 248-250
- Long Beach, California: site of study on growth rate, distribution and population density of northern quahog, 68-94
- Measurement: of salmon using an old and unfamiliar method, 247
- Method, new: of evaluating fish species composition in limnetic areas of reservoirs, 209-214
- Method, old: of measuring salmon, 247
- Minnow, fathead: occurrence in Oregon, 155-156
- Modification: of Clover deer trap, 242-244
- Mollusks: range extension California, 152-155; size records in California, 152-155
- Molt: of California condor as field study aid, 144-148
- Monterey Bay: occurrence of North Pacific frostfish, 149-152; occurrence of prowlfish, 60-62; occurrence of parakeet auklet, 158; occurrence of pugheadedness in longspine combfish, 160-162
- Mortality: of two different sizes of silver salmon stocked in Lake Berryessa, 127-132
- Observations: on food habits of leopard sharks and brown smoothhounds, 95-103
- Occurrence: of fathead minnow in Oregon, 155-156; of North Pacific froshfish in Monterey Bay, 149-152; of parakeet auklet at Monterey Bay, 158; of prowlfish in Monterey Bay, 60-62; of yellow snake eel off southern California, 157-158
- Otoliths: used as a means to calculate growth rate of northern anchovy in southern California waters, 116-126
- Otoliths, anomalous: from the northern anchovy, 235-236
- Parasites: of fishes from Sacramento-San Joaquin Delta, 201-208
- Parasites, external: of California inshore sharks, 228-232
- Plumage: of California condor as field study aid, 144-148
- Polymorphism: in populations of *Sceloporus occidentalis* in Santa Barbara County, 253-254

- Population density: of northern quahog in Long Beach, California, 68-81
 Population study: of black bear in northern California, 4-25
 Populations: of *Sceloporus occidentalis* in Santa Barbara County exhibiting polymorphism, 253-254
 Prowfish: occurrence in Monterey Bay, 60-62
 Pugheadedness: in longspine combfish, 160-162
 Quahog, northern: growth rate, 68-94; distribution, 68-94; population density, 68-94
 Range extension: for *Terebra pedroana* in California, 152-155
 Range extension, depth: of the yellow snake eel, 157-158
 Range extension, latitudinal: for yellow snake eel, 156-157; for spotted snake eel, 156-157
 Ratfish: note on ecology of, 47-53
 Rays, bat: an unusual aggregation of, 159
 Record, first: of *Anthias gordenis*, 111-112
 Record, further: of Little Kern golden trout in Little Kern River Basin, 248-250
 Records, size: for mollusks in California, 152-155
 Reservoirs: evaluating fish species composition in limnetic areas by fish trapping, 200-214
 Reviews: A river never sleeps, 258; Fisherman's spring, 260; Fly casting with Lefty Kreh, 256; Endemism in fishes of the Clear Lake region of central California, 63; Freshwater ecology, 255; How to tie freshwater flies, 256; Marine molluscan genera of western North America: An illustrated key, 259; Return to the river, 257; Seashore life of Puget Sound, the Strait of Georgia, and the San Juan Archipelago, 259-260; The blackbass in America and overseas, 260; The carnivores, 255-256; The shore fishes of Hawaii, 258-259; The trout and the stream, 257-258; Trout magic, 257; Western trout fly tying manual, 63; Wilderness fishing for salmon and steelhead, 257
 Richardson Bay: a list of fishes collected, 104-106
 Sacramento River: site of study analyzing blood and serum of adult striped bass, 245-246
 Sacramento-San Joaquin Delta: site of study on fish parasites, 201-208
 Sacramento-San Joaquin Estuary: site of study on food habits of *Neomysis mercedis*, 35-46
 Salmon: an old and unfamiliar method of measuring, 247
 Salmon, silver: mortality rates, 127-132; growth rates of two different sizes, 127-132; cost of two different sizes, 127-132
 San Francisco Bay: changes in species composition of sharks, 106-111
 Sea lion, Steller, extrauterine fetus in, 233-234
 Sharks: changes in species composition in south San Francisco Bay, 106-111
 Sharks, brown smoothhound: food habits, 95-103
 Sharks, inshore: external parasites collected from, 228-232
 Sharks, leopard: food habits, 95-103
 Snake eel, yellow: depth range extension, 157-158; latitudinal range extension, 156-157; occurrence off southern California, 157-158
 Snake eel, spotted: latitudinal range extension, 156-157
 Species composition: changes in shark species composition in south San Francisco Bay, 106-111; of fish in limnetic areas of reservoirs as determined by fish trapping, 200-214
 Status: of Rocky Mountain elk in Kern County, 1974, 239-241
 Status, breeding: of tricolored blackbird, 1960-1972, 166-180
 Study aids: used in California condor identification, 144-148
 Trap, corral: Stewart modified type used in trapping deer, 237-239
 Trap, deer: modification of Clover deer trap, 242-244
 Trapping: a new method of evaluating fish species composition in limnetic areas of reservoirs, 200-214
 Trout, golden: further record of Little Kern golden trout in Little Kern River Basin, 248-250
 Turbot, diamond: larvae of, 26-34





